

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF TEXAS

This label for Onager expires and must not be distributed or used in accordance with this SLN registration after December 31, 2023.



EMULSIFIABLE CONCENTRATE FOR AGRICULTURAL USE ONLY

ACTIVE INGREDIENT: Hexythiazox	% By Wt.
<i>trans</i> -5-(4-Chlorophenyl)-N-cyclohexyl-4-methyl-2-oxothiazolidine-3-carboxamide	11.8%
OTHER INGREDIENTS	88.2%
	TOTAL 100.0%

Onager contains 1.0 lb. active ingredient per gallon
Contains Petroleum Distillate

KEEP OUT OF REACH OF CHILDREN CAUTION

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable restrictions, precautions, worker protection standard requirements, and directions for use on the EPA label must be followed.
- This labeling must be in the possession of the user at the time of pesticide application.

CROP	PEST	RATE Oz./acre	COMMENTS
Potatoes*	Two spotted spider mite, Potato Psyllid	16-24	Ground Application: Apply at recommended rates Air Application: Apply a minimum of 5 gallons finished spray per acre. Chemigation: see details under Chemigation Systems on Section 3 label.
	<ul style="list-style-type: none"> Do not apply within 21 days of harvest Do not apply more than a total of 24 oz. formulated product per acre per calendar year. Do not make more than one application per calendar year. *May be applied in eastern Texas: TX: E of Rt. 283 and SE of Rt. 377		

RESTRICTIONS ON ALL CROPS

- Do not graze or feed livestock on cover crops growing in treated areas.
- Onager is an emulsifiable concentrate to be diluted with water for application in commercial plantings only.
- Apply Onager prior to adult mite build up. Onager will not control adult spider mites. Use higher rates on moderate to high mite infestations or for larger plants with a dense canopy. If adult mites are present in medium to high populations, better results may be obtained using Onager in combination with a registered contact adulticide. The use of less than label rates is not recommended with Onager.
- Do not make more than one application of Onager or any other Hexythiazox product to the same crop per year.
- Do not plant rotational crops other than those on the Section 3 label within 120 days of this application.